

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Anti-Hu TCR gamma/delta PE-Cy(TM)7, Clone: [11F2], PE/Cy7, Monoclonal EXB-T7-912-T100

| | |
|--------------------------|---|
| Artikelname | Anti-Hu TCR gamma/delta PE-Cy(TM)7, Clone: [11F2], PE/Cy7, Monoclonal |
| Artikelnummer | EXB-T7-912-T100 |
| Hersteller Artikelnummer | T7-912-T100 |
| Alternativnummer | EXB-T7-912-T100 |
| Hersteller | EXBIO |
| Kategorie | Antikörper |
| Applikation | FC |
| Spezies Reaktivität | Human |
| Immunogen | Purified CD3 antigen complex |
| Konjugation | PE/Cy7 |
| Produktbeschreibung | The antigen-specific T cell receptor (TCR) is composed of either alpha and beta subunit, or gamma and delta subunit. Majority of T cells present in the blood, lymph and secondary lymphoid organs express TCR alpha/beta heterodimers, whereas the T cell... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [11F2] |
| Isotyp | Mouse IgG1 |
| Puffer | Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide |

| | |
|------------------------|--|
| Target-Kategorie | TCR gamma/delta |
| Antibody Type | Monoclonal Antibody |
| Application Verdünnung | Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests. |
| Anwendungsbeschreibung | Flow cytometry: The reagent is designed for analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests. |