

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Mouse anti Human FAS/APO-1 (CD95), IgG1, Clone: [3.22], FITC, Monoclonal NMB-0952

| | |
|--------------------------|---|
| Artikelname | Mouse anti Human FAS/APO-1 (CD95), IgG1, Clone: [3.22], FITC, Monoclonal |
| Artikelnummer | NMB-0952 |
| Hersteller Artikelnummer | 0952 |
| Alternativnummer | NMB-0952 |
| Hersteller | NordicMubio |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC |
| Spezies Reaktivität | Human |
| Konjugation | FITC |
| Produktbeschreibung | Activation of either the 55-kD tumor necrosis factor receptor (TNF-R1) or CD95 (Fas/Apo-1) causes apoptosis of cells and liver failure in mice, and has been associated with human liver disorders. The aim of this study was first to clarify the associa... |
| Klonalität | Monoclonal |
| Konzentration | See vial for concentration |
| Klon-Bezeichnung | [3.22] |
| Isotyp | IgG1 |

| | |
|------------------------|---|
| UniProt | P25445 |
| Puffer | Provided as solution in phosphate buffered saline with 0.08% sodium azide and 0.2% carrier protein |
| Reinheit | Protein A/G Chromatography |
| Formel | Provided as solution in phosphate buffered saline with 0.08% sodium azide and 0.2% carrier protein |
| Anwendungsbeschreibung | PBMC: Add 10 µl of MAB/10 6 PBMC in 100 µl PBS. Mix gently and incubate for 15 minutes at 2 to 8C. Wash twice with PBS and analyze or fix with 0.5% v/v of paraformaldehyde in PBS and analyze. WHOLE BLOOD: Add 10 µl of MAB/100 µl of whole blood. Mix gently |