

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Mouse anti Human FAS/APO-1 (CD95), conjugated with PE, IgG1, Clone: [3.22], RPE, Monoclonal NMB-0954

| | |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname | Mouse anti Human FAS/APO-1 (CD95), conjugated with PE, IgG1, Clone: [3.22], RPE, Monoclonal |
| Artikelnummer | NMB-0954 |
| Hersteller Artikelnummer | 0954 |
| Alternativnummer | NMB-0954 |
| Hersteller | NordicMubio |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC |
| Spezies Reaktivität | Human |
| Immunogen | Full length CD95 protein |
| Konjugation | RPE |
| Produktbeschreibung | Activation of either the 55-kD tumor necrosis factor receptor (TNF-R1) or CD95 (Fas/Apo-1) causes apoptosis of cells and liver failure in mice, and has been associated with human liver disorders. The aim of this study was first to clarify the associa... |
| Klonalität | Monoclonal |
| Konzentration | Titered for flow cytometry |
| Klon-Bezeichnung | [3.22] |

| | |
|------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Isotyp | IgG1 |
| UniProt | P25445 |
| Puffer | Provided as solution in phosphate buffered saline with 0.08% sodium azide |
| Reinheit | Protein A/G Chromatography |
| Formel | Provided as solution in phosphate buffered saline with 0.08% sodium azide |
| Anwendungsbeschreibung | PBMC: Add 10 µl of MAB/10 ⁶ PBMC in 100 µl PBS. Mix gently and incubate for 15 minutes at 2 to 8°C. Wash twice with PBS and analyze or fix with 0.5% v/v of paraformaldehyde in PBS and analyze. WHOLE BLOOD: Add 10 µl of MAB/100 µl of whole blood. Mix gently |