

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Mouse anti double-stranded RNA (J2) NMB-10010200

| | |
|--------------------------|--|
| Artikelname | Mouse anti double-stranded RNA (J2) |
| Artikelnummer | NMB-10010200 |
| Hersteller Artikelnummer | 10010200 |
| Alternativnummer | NMB-10010200 |
| Hersteller | NordicMubio |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | DOT, ELISA, FC, IAC, IB, ICC, IHC |
| Produktbeschreibung | Over the past decade our double-stranded RNA (dsRNA)antibodies have been used extensively to detect and characterise plant and animal viruses with dsRNA genomes or intermediates. In addition, the anti-dsRNA antibodies can be used as a diagnostic tool... |
| Klonalität | Monoclonal |
| Konzentration | Concentration after reconstitution: 1.00 mg/ml as determined by A280 nm (A280 nm = 1.47 corresponds to 1 mg/ml antibody). |
| Klon-Bezeichnung | J2 |
| Isotyp | IgG2a kappa |
| Puffer | Mouse monoclonal antibody J2 recognises double-stranded RNA (dsRNA) provided that the length of the helix is greater than or equal to 40 bp. dsRNA-recognition is independent of the sequence and nucleotide composition of the antigen. All naturally occurri |

| | |
|------------------------|---|
| Quelle | Female DBA/2 mice were injected intraperitoneally with a mixture of 50 ug L-dsRNA and 75 ug methylated bovine serum albumin, emulsified in complete Freund's adjuvant. After several boosts spleen cells were fused with Sp2/0-Ag14 myeloma cells to generate t |
| Reinheit | Gel electrophoretically pure IgG antibody. |
| Formel | The lyophilised sample should be reconstituted with 200 µl sterile distilled water. The mAb will then be in PBS without any stabilisers at a concentration of 1 mgr/ml. As a result of the lyophilisation procedure, the reconstituted antibody may contain sm |
| Anwendungsbeschreibung | Mouse monoclonal antibody J2 can be used for ELISA, dsRNA-immunoblotting, immunoaffinity chromatography and in certain systems also for immunohistochemistry (see references). The optimum working dilution of the antibody for any specific application shoul |