

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

EpCAM/CD326, IgG1, Clone: [HEA125], Mouse, Monoclonal NMB-MUB2061P

| | |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname | EpCAM/CD326, IgG1, Clone: [HEA125], Mouse, Monoclonal |
| Artikelnummer | NMB-MUB2061P |
| Hersteller Artikelnummer | MUB2061P |
| Alternativnummer | NMB-MUB2061P |
| Hersteller | NordicMubio |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | ELISA, FC, IF, IHC |
| Spezies Reaktivität | Human |
| Immunogen | Viable HT-29 human colon carcinoma cell line |
| Produktbeschreibung | Ep-CAM is a 34 kD glycoprotein and can be detected at the basolateral membrane of the majority of epithelial tissues. It is intricately linked with the cadherin-catenin complex and hence the fundamental WNT pathway responsible for intracellular signa... |
| Klonalität | Monoclonal |
| Konzentration | 1 mg/ml |
| Klon-Bezeichnung | [HEA125] |
| Isotyp | IgG1 |
| UniProt | P16422 |

| | |
|------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Puffer | Each vial contains 100µg 1mg/ml purified monoclonal antibody in PBS containing 0.09% sodium azide. |
| Quelle | HEA125 is a mouse monoclonal IgG1 antibody derived by fusion of Ag8.653 mouse myeloma cells with splenocytes from a BALB/c mouse cells immunized with the HT-29 colon carcinoma cell line. |
| Reinheit | ProtG affinity chromatography |
| Anwendungsbeschreibung | The HEA125 antibody is suitable for the detection of human EpCAM by Western blotting (under reducing and non-reducing conditions), flow cytometry and immunocytochemistry. The antibody is also suitable for the detection of human EpCAM by immunohistochemis |