

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Mouse anti Mitotic cells, IgM, Clone: [8B3G], Monoclonal NMB-MUB1200P

| | |
|--------------------------|---|
| Artikelname | Mouse anti Mitotic cells, IgM, Clone: [8B3G], Monoclonal |
| Artikelnummer | NMB-MUB1200P |
| Hersteller Artikelnummer | MUB1200P |
| Alternativnummer | NMB-MUB1200P |
| Hersteller | NordicMubio |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, ICC |
| Spezies Reaktivität | Human, Zebrafish |
| Produktbeschreibung | The life cycle of a eukaryotic cell consists of various phases, two of which can morphologically and biochemically be identified. Firstly, during mitosis (M-phase), in which the cell divides into two identical daughter cells, chromosome condensation ... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [8B3G] |
| Isotyp | IgM |
| Puffer | Each vial contains 100 ul 1 mg/ml purified monoclonal antibody in PBS containing 0.09% sodium azide. The product may contain some contaminating proteins next to the IgM as a result of ammonium sulphate precipitation being used for antibody purification. |

| | |
|------------------------|--|
| Quelle | 8B3G is a mouse monoclonal IgM antibody derived by fusion of SP2/0-Ag14 Mouse myeloma cells with spleen cells from a mouse immunized with a total cell lysate of the human bladder carcinoma cell line T24. |
| Reinheit | Ammonium sulphate precipitation |
| Anwendungsbeschreibung | 8B3G can be used for flow cytometric analyses and immunocytochemistry. 8B3G is not suitable for immunoblotting. Optimal antibody dilution should be determined by titration, recommended range is 1:50 - 1:100 for flow cytometry, and for immunocytochemistry |