

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat anti Bovine IgG (heavy and light chains), conjugated with Biotin, Clone: [Polyclonal], Monoclonal NMB-GAB/IGG(H+L)/BIO

| | |
|--------------------------|--|
| Artikelname | Goat anti Bovine IgG (heavy and light chains), conjugated with Biotin, Clone: [Polyclonal], Monoclonal |
| Artikelnummer | NMB-GAB/IGG(H+L)/BIO |
| Hersteller Artikelnummer | GAB/IgG(H+L)/Bio |
| Alternativnummer | NMB-GAB/IGG(H+L)/BIO |
| Hersteller | NordicMubio |
| Wirt | Goat |
| Kategorie | Antikörper |
| Konjugation | Biotin |
| Produktbeschreibung | The reactivity of the antiserum is directed to the Fc and Fab subunits of the IgG molecule. It includes a certain degree of reactivity with other immunoglobulins via the common Fab portion. The reactivity against Fab of IgG is less pronounced than in... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [Polyclonal] |
| Puffer | Biotin-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added It is reconstituted by adding 1 ml sterile distilled water, spun down to remove insoluble particles, divided into s |

| | |
|------------------------|--|
| Quelle | Purified normal IgG isolated from pooled bovine serum. Freund's complete adjuvant is used in the first step of the immunization procedure. |
| Formel | Biotin-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added |
| Anwendungsbeschreibung | ELISA, Immunocytochemistry, Immunohistochemistry (paraffin), Dot blot, Immunoblotting. |