

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat anti Chicken IgA (Fc specific), conjugated with FITC, Clone: [Polyclonal], Monoclonal NMB-GACH/IGA(FC)/FITC

| | |
|--------------------------|--|
| Artikelname | Goat anti Chicken IgA (Fc specific), conjugated with FITC, Clone: [Polyclonal], Monoclonal |
| Artikelnummer | NMB-GACH/IGA(FC)/FITC |
| Hersteller Artikelnummer | GACH/IgA(Fc)/FITC |
| Alternativnummer | NMB-GACH/IGA(FC)/FITC |
| Hersteller | NordicMubio |
| Wirt | Goat |
| Kategorie | Antikörper |
| Konjugation | FITC |
| Produktbeschreibung | The reactivity of the antiserum is directed to the Fc subunit of the IgA molecule, which expresses strict (class) specificity. In immunoelectrophoresis and radial in immunodiffusion using various antiserum concentrations against chicken serum, a sing... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [Polyclonal] |
| Puffer | FITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1 ml sterile distilled water, spun dow |
| Quelle | Purified IgA isolated from chicken serum. Freund's complete adjuvant is used in the first step of the immunization procedure. |

| | |
|------------------------|---|
| Formel | FITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. |
| Anwendungsbeschreibung | ELISA, Immunocytochemistry, Immunohistochemistry (frozen), (In)direct immunofluorescence. |