

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat anti Guinea Pig IgG (subclass specific), Clone: [Polyclonal], Monoclonal NMB-GAGP/IGG1

| | |
|--------------------------|--|
| Artikelname | Goat anti Guinea Pig IgG (subclass specific), Clone: [Polyclonal], Monoclonal |
| Artikelnummer | NMB-GAGP/IGG1 |
| Hersteller Artikelnummer | GAGp/IgG1 |
| Alternativnummer | NMB-GAGP/IGG1 |
| Hersteller | NordicMubio |
| Wirt | Goat |
| Kategorie | Antikörper |
| Produktbeschreibung | The reactivity of the antiserum is directed to the Fc subunit of subclass IgG1. It does not react with other IgG2, IgG/Fab fragments, IgM and IgA or any non-Ig protein in guinea pig serum, as tested by immunoelectrophoresis and double radial immunodi... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [Polyclonal] |
| Puffer | Delipidated, heat inactivated, lyophilized whole antiserum. No preservative added, as it may interfere with the antibody activity. Total protein and IgG concentration in the antiserum are comparable to those of pooled normal goat serum. No foreign protei |
| Quelle | Pools of purified normal IgG1 isolated from pooled guinea pig serum. Freund's complete adjuvant is used in the first step of the immunization procedure. |

| | |
|------------------------|--|
| Formel | Delipidated, heat inactivated, lyophilized whole antiserum. No preservative added, as it may interfere with the antibody activity. Total protein and IgG concentration in the antiserum are comparable to those of pooled normal goat serum. No foreign protei |
| Anwendungsbeschreibung | Precipitation assays. In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and |