

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat anti Guinea Pig Serum proteins, Clone: [Polyclonal], Monoclonal NMB-GAGP/TSP

| | |
|--------------------------|---|
| Artikelname | Goat anti Guinea Pig Serum proteins, Clone: [Polyclonal], Monoclonal |
| Artikelnummer | NMB-GAGP/TSP |
| Hersteller Artikelnummer | GAGp/TSP |
| Alternativnummer | NMB-GAGP/TSP |
| Hersteller | NordicMubio |
| Wirt | Goat |
| Kategorie | Antikörper |
| Produktbeschreibung | In immunoelectrophoresis against pooled serum and enriched serum proteins fractions precipitation can be observed of not less than 18 individual proteins components. In precipitating techniques as immunoelectrophoresis and radial immunodiffusion (Ouc... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [Polyclonal] |
| Puffer | Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. Reconstitute the lyophilized |
| Quelle | Pooled whole guinea pig serum and partly purified serum fractions. Freund's complete adjuvant is used in the first step of the immunization procedure. |

| | |
|------------------------|---|
| Formel | Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. |
| Anwendungsbeschreibung | Precipitation assays. In immunoelectrophoresis use 2 µl serum or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in 3 mm diameter center well and 2 µl serum samples (neat |