

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat anti Human C4, Clone: [Polyclonal], Monoclonal NMB-GAHU/C4

| | |
|--------------------------|--|
| Artikelname | Goat anti Human C4, Clone: [Polyclonal], Monoclonal |
| Artikelnummer | NMB-GAHU/C4 |
| Hersteller Artikelnummer | GAHu/C4 |
| Alternativnummer | NMB-GAHU/C4 |
| Hersteller | NordicMubio |
| Wirt | Goat |
| Kategorie | Antikörper |
| Produktbeschreibung | The reactivity of the antiserum is restricted to C4. In immunoelectrophoresis and radial immunodiffusion (Ouchterlony), using various antiserum concentrations against fresh normal human plasma a single precipitin line is obtained which shows a reacti... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [Polyclonal] |
| UniProt | POCOL5 |
| Puffer | Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. Reconstitute the lyophilized a |
| Quelle | C4 and C2 represent substrates for activated C1. Both components are present in serum in an inert form. C4 is composed of 3 polypeptide chains of three different types, , , . One of the chains is cleaved by the action of activated C1, resulting in activa |

| | |
|------------------------|---|
| Formel | Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. |
| Anwendungsbeschreibung | Precipitation assays. Dilutions may be prepared by adding phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect |