

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat anti Human IgD (Fc specific), conjugated with FITC, Clone: [Polyclonal], Monoclonal NMB-SR/GAHU/IGD(FC)/FITC

| | |
|--------------------------|--|
| Artikelname | Goat anti Human IgD (Fc specific), conjugated with FITC, Clone: [Polyclonal], Monoclonal |
| Artikelnummer | NMB-SR/GAHU/IGD(FC)/FITC |
| Hersteller Artikelnummer | SR/GAHu/IgD(Fc)/FITC |
| Alternativnummer | NMB-SR/GAHU/IGD(FC)/FITC |
| Hersteller | NordicMubio |
| Wirt | Goat |
| Kategorie | Antikörper |
| Konjugation | FITC |
| Produktbeschreibung | The reactivity of the antiserum is directed to the Fc subunit of the IgD molecule, which expresses strict (class) specificity. In immunoelectrophoresis and radial in immunodiffusion using various antiserum concentrations against human serum, a single... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [Polyclonal] |
| Puffer | FITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1 ml sterile distilled water, spun dow |

| | |
|------------------------|---|
| Quelle | Pools of purified homogenous IgD isolated from human serum. Freund's complete adjuvant is used in the first step of the immunization procedure. |
| Formel | FITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. |
| Anwendungsbeschreibung | ELISA, Immunocytochemistry, Immunohistochemistry (frozen), (In)direct immunofluorescence. |