

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat anti Human J chain of dimeric IgA, Clone: [Polyclonal], Monoclonal NMB-GAHU/J

| | |
|--------------------------|--|
| Artikelname | Goat anti Human J chain of dimeric IgA, Clone: [Polyclonal], Monoclonal |
| Artikelnummer | NMB-GAHU/J |
| Hersteller Artikelnummer | GAHu/J |
| Alternativnummer | NMB-GAHU/J |
| Hersteller | NordicMubio |
| Wirt | Goat |
| Kategorie | Antikörper |
| Applikation | IP |
| Spezies Reaktivität | Human |
| Produktbeschreibung | The reactivity of the antiserum is restricted to J chain as tested in immunoelectrophoresis and radial immunodiffusion. The antiserum shows a single precipitation reaction with totally reduced and alkylated polyclonal and monoclonal polymeric IgA and... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [Polyclonal] |
| Puffer | Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. Reconstitute the lyophilized a |

| | |
|------------------------|--|
| Quelle | Human J chain is a polypeptide folded within the polymeric structure of the immunoglobulin. J chain isolated from human polymeric IgA and IgM are identical by criteria of composition, peptide maps and antigenicity. Human J chain has been established as d |
| Anwendungsbeschreibung | Precipitation assays. In immunoelectrophoresis use 2 μ l serum or equivalent against 120 μ l antiserum. In double radial immunodiffusion use a rosette arrangement with 10 μ l antiserum in 3 mm diameter centre well and 2 μ l serum samples (neat and serially d |