

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat anti Human thyroxine binding globulin, Clone: [Polyclonal], Monoclonal NMB-GAHU/TBG

| | |
|--------------------------|--|
| Artikelname | Goat anti Human thyroxine binding globulin, Clone: [Polyclonal], Monoclonal |
| Artikelnummer | NMB-GAHU/TBG |
| Hersteller Artikelnummer | GAHu/TBG |
| Alternativnummer | NMB-GAHU/TBG |
| Hersteller | NordicMubio |
| Wirt | Goat |
| Kategorie | Antikörper |
| Produktbeschreibung | The reactivity of the antiserum is restricted to thyroxine binding globulin. In immunoelectrophoresis and radial immunodiffusion (Ouchterlony), using various antiserum concentrations against fresh normal human plasma a single precipitin line is obtai... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [Polyclonal] |
| UniProt | P05543 |
| Puffer | Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. Reconstitute the lyophilized an |

| | |
|------------------------|---|
| Quelle | Thyroxine binding is a protein with a molecular weight of 55,700 kD. The protein is binding thyroxine as well as triiodothyronine. In immunoelectrophoresis it has a alpha mobility. The mean concentration in normal human serum is about 40 µg/ml. Elevated |
| Formel | Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added. |
| Anwendungsbeschreibung | Precipitation assays. Dilutions may be prepared by adding phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect |