

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Goat anti Monkey IgM (Fc specific), conjugated with Biotin, Clone: [Polyclonal], Monoclonal NMB-GAMON/IGM(FC)/BIO**

|                          |  |
|--------------------------|--|
| Artikelname              | Goat anti Monkey IgM (Fc specific), conjugated with Biotin, Clone: [Polyclonal], Monoclonal  |
| Artikelnummer            | NMB-GAMON/IGM(FC)/BIO  |
| Hersteller Artikelnummer | GAMon/IgM(Fc)/Bio  |
| Alternativnummer         | NMB-GAMON/IGM(FC)/BIO  |
| Hersteller               | NordicMubio  |
| Wirt                     | Goat   |
| Kategorie                | Antikörper   |
| Konjugation              | Biotin   |
| Produktbeschreibung      | The reactivity of the antiserum is directed to the Fc subunit of the IgM molecule which expresses strict isotypic (class) specificity. It does not react with IgG, IgA and IgG/Fab or any non-Ig protein in monkey serum, as tested by immunoelectrophores...  |
| Klonalität               | Monoclonal   |
| Klon-Bezeichnung         | [Polyclonal]   |
| Puffer                   | Biotin-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1 ml sterile distilled water, spun d |
| Quelle                   | Purified IgM isolated from Rhesus monkey serum. Freund's complete adjuvant is used in the first step of the immunization procedure.  |

|                        |   |
|------------------------|---|
| Formel                 | Biotin-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. |
| Anwendungsbeschreibung | ELISA, Immunocytochemistry, Immunohistochemistry (paraffin), Dot blot, Immunoblotting.  |