

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Goat anti Mouse IgG1 IgG2a IgG2b IgG3 (Fc specific), Clone: [Polyclonal], Monoclonal NMB-GAM/IGG(FC)**

|                          |  |
|--------------------------|--|
| Artikelname              | Goat anti Mouse IgG1 IgG2a IgG2b IgG3 (Fc specific), Clone: [Polyclonal], Monoclonal   |
| Artikelnummer            | NMB-GAM/IGG(FC)  |
| Hersteller Artikelnummer | GAM/IgG(Fc)  |
| Alternativnummer         | NMB-GAM/IGG(FC)  |
| Hersteller               | NordicMubio  |
| Wirt                     | Goat   |
| Kategorie                | Antikörper   |
| Produktbeschreibung      | The reactivity of the antiserum is directed to the Fc subunit of the IgG molecule. The antiserum contains antibodies to subclass-specific determinants as well as to determinants shared by two or more subclasses of IgG. In immunoelectrophoresis and do...  |
| Klonalität               | Monoclonal   |
| Klon-Bezeichnung         | [Polyclonal]   |
| Puffer                   | Delipidated, heat inactivated, lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. Total protein and IgG concentration in the antiserum are comparable to those of pooled goat serum. No foreign proteins a |
| Quelle                   | Purified normal IgG, including all known subclasses, isolated from pooled mouse serum. Freund's complete adjuvant is used in the first step of the immunization procedure.   |

|                        |  |
|------------------------|--|
| Formel                 | Delipidated, heat inactivated, lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. Total protein and IgG concentration in the antiserum are comparable to those of pooled goat serum. No foreign proteins a |
| Anwendungsbeschreibung | Precipitation assays. In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and  |