

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Goat anti Rabbit IgG IgA IgM (heavy and light chains), conjugated with Biotin, Clone: [Polyclonal], Monoclonal NMB-GAR/IG/BIO

| | |
|--------------------------|--|
| Artikelname | Goat anti Rabbit IgG IgA IgM (heavy and light chains), conjugated with Biotin, Clone: [Polyclonal], Monoclonal |
| Artikelnummer | NMB-GAR/IG/BIO |
| Hersteller Artikelnummer | GAR/Ig/Bio |
| Alternativnummer | NMB-GAR/IG/BIO |
| Hersteller | NordicMubio |
| Wirt | Goat |
| Kategorie | Antikörper |
| Konjugation | Biotin |
| Produktbeschreibung | The reactivity of the antiserum is directed to the Fc and Fab subunits of the major immunoglobulin es. It does not react with any non-Ig protein in rabbit serum, as tested by immunoelectrophoresis and double radial immunodiffusion. In immunocytochemi... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [Polyclonal] |
| Puffer | Biotin-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1 ml sterile distilled water, spun d |

| | |
|------------------------|---|
| Quelle | Purified normal IgG, IgA and IgM isolated from pooled rabbit serum. Freund's complete adjuvant is used in the first step of the immunization procedure. |
| Formel | Biotin-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. |
| Anwendungsbeschreibung | ELISA, Immunocytochemistry, Immunohistochemistry (paraffin), Dot blot, Immunoblotting. |