

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Goat anti Rat IgG2c (subclass specific), conjugated with TRITC, Clone: [Polyclonal], Monoclonal NMB-GARA/IGG2C/TRITC**

|                          |  |
|--------------------------|--|
| Artikelname              | Goat anti Rat IgG2c (subclass specific), conjugated with TRITC, Clone: [Polyclonal], Monoclonal  |
| Artikelnummer            | NMB-GARA/IGG2C/TRITC   |
| Hersteller Artikelnummer | GARa/IgG2c/TRITC   |
| Alternativnummer         | NMB-GARA/IGG2C/TRITC   |
| Hersteller               | NordicMubio  |
| Wirt                     | Goat   |
| Kategorie                | Antikörper   |
| Konjugation              | TRITC  |
| Produktbeschreibung      | The reactivity of the antiserum is directed to the subclass IgG2c. It does not react with other subclasses of IgG, IgG/Fab fragments, IgM and IgA or any non-Ig protein in rat serum, as tested by immunoelectrophoresis and double radial immunodiffusion...  |
| Klonalität               | Monoclonal   |
| Klon-Bezeichnung         | [Polyclonal]   |
| Puffer                   | TRITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1 ml sterile distilled water, spun do |

|                        |  |
|------------------------|--|
| Quelle                 | Pools of purified homogenous IgG2c isolated from rat sera. Freund's complete adjuvant is used in the first step of the immunization procedure.   |
| Formel                 | TRITC-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. |
| Anwendungsbeschreibung | ELISA, Immunocytochemistry, Immunohistochemistry (frozen), (In)direct immunofluorescence.  |