

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Mouse anti Human IgA2 (A2(m)2 allotype specific), IgG1, Clone: [NI 194-3 A89-040], Monoclonal NMB-MAHU/IGA2(M)2**

Artikelname	Mouse anti Human IgA2 (A2(m)2 allotype specific), IgG1, Clone: [NI 194-3 A89-040], Monoclonal
Artikelnummer	NMB-MAHU/IGA2(M)2
Hersteller Artikelnummer	MAHu/IgA2(m)2
Alternativnummer	NMB-MAHU/IGA2(M)2
Hersteller	NordicMubio
Wirt	Mouse
Kategorie	Antikörper
Applikation	ELISA, ICC, IHC, WB
Spezies Reaktivität	Human
Produktbeschreibung	The reactivity of the antiserum is restricted to the IgA2(m)2 molecule as tested in haemagglutination, haemagglutination inhibition, direct binding enzyme immunoassay, competitive inhibition enzyme immunoassay, immunoblotting, immunoprecipitation, la...
Klonalität	Monoclonal
Klon-Bezeichnung	[NI 194-3 A89-040]
Isotyp	IgG1
Puffer	Delipidated, heat inactivated, lyophilized, stable whole ascites IgG concentration is 1 mg/ml. No foreign proteins added. Reconstitute the lyophilized ascites by adding 0.5 ml sterile distilled water.

Quelle	Highly purified monoclonal IgA2(m)2 isolated from human serum.
Formel	Delipidated, heat inactivated, lyophilized, stable whole ascites IgG concentration is 1 mg/ml. No foreign proteins added.
Anwendungsbeschreibung	Indirect immunofluorescence,ELISA. The optimum working dilution is an assay-related characteristic and should always be determined by titration. For histochemical use optimum dilutions are mostly from 1:50 to 1:200, in ELISA from 1:200 upwards, in Wester