

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Mouse anti Myeloperoxidase-C2 (MPO-C2), conjugated to FITC, IgG1, Clone: [8E6], Monoclonal NMB-GM-4192

| | |
|--------------------------|--|
| Artikelname | Mouse anti Myeloperoxidase-C2 (MPO-C2), conjugated to FITC, IgG1, Clone: [8E6], Monoclonal |
| Artikelnummer | NMB-GM-4192 |
| Hersteller Artikelnummer | GM-4192 |
| Alternativnummer | NMB-GM-4192 |
| Hersteller | NordicMubio |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC |
| Spezies Reaktivität | Human |
| Konjugation | FITC |
| Produktbeschreibung | Myeloperoxidase (MPO) is a glycoprotein present in the azurophil (primary) granules of myeloid cells, which appears in the myeloblast stage of myeloid cell differentiation. MPO is the most common functional protein of myeloid cells and is involved in ... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [8E6] |
| Isotyp | IgG1 |
| UniProt | P05164 |

| | |
|------------------------|---|
| Puffer | 2 ml of FITC-conjugated anti Myeloperoxidase-C2 (clone 8E6) in PBS pH 7.2, 1% BSA, and 0.05% NaN ₃ , approximately 100 tests. |
| Reinheit | Purified by Chromatography |
| Formel | PBS pH 7.2, 1% BSA, 0.05% NaN ₃ |
| Anwendungsbeschreibung | Permeabilization and Staining Procedure - In combination with our Permeabilization Kit FIX&PERM (Cat. No. GAS-002) intracellular MPO can be easily stained in cell suspensions. - For each sample to be analyzed add 50 µl of whole blood, bone marrow or mon |