

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Rabbit anti Goat IgM (heavy and light chains), Clone: [Polyclonal], Monoclonal NMB-RAG/IGM(H+L)**

Artikelname	Rabbit anti Goat IgM (heavy and light chains), Clone: [Polyclonal], Monoclonal
Artikelnummer	NMB-RAG/IGM(H+L)
Hersteller Artikelnummer	RAG/IgM(H+L)
Alternativnummer	NMB-RAG/IGM(H+L)
Hersteller	NordicMubio
Wirt	Rabbit
Kategorie	Antikörper
Produktbeschreibung	The reactivity of the antiserum is directed to the Fc and Fab subunits of the IgM molecule. It includes a certain degree of reactivity with other immunoglobulins via the common Fab portion. It does not react with any non-Ig protein in goat serum, as ...
Klonalität	Monoclonal
Klon-Bezeichnung	[Polyclonal]
Puffer	Delipidated, heat inactivated lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity.No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled rabbit se
Quelle	Purified normal IgM isolated from pooled goat serum. Freund's complete adjuvant is used in the first step of the immunization procedure.

Formel	Delipidated, heat inactivated lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity.No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled rabbit se
Anwendungsbeschreibung	Precipitation assays. In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and