

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Rabbit anti Human C2, Clone: [Polyclonal], Monoclonal NMB-RAHU/C2

|                          |   |
|--------------------------|---|
| Artikelname              | Rabbit anti Human C2, Clone: [Polyclonal], Monoclonal   |
| Artikelnummer            | NMB-RAHU/C2   |
| Hersteller Artikelnummer | RAHu/C2   |
| Alternativnummer         | NMB-RAHU/C2   |
| Hersteller               | NordicMubio   |
| Wirt                     | Rabbit  |
| Kategorie                | Antikörper  |
| Produktbeschreibung      | The reactivity of the antiserum is restricted to native and activated C2. In immunoelectrophoresis and radial immunodiffusion, using various antiserum concentrations against human plasma a single precipitin line is obtained and shows a reaction of fu... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [Polyclonal]  |
| UniProt                  | <a href="#">P06681</a>  |
| Puffer                   | Delipidated, heat inactivated, lyophilized, stable IgG fraction. No preservative added. The IgG concentration is 4 mg/ml in PBS-EDTA. No foreign proteins added. Reconstitute the lyophilized antiserum by adding 1 ml sterile distilled water.               |

|                        |  |
|------------------------|--|
| Quelle                 | The C2 component of human complement is a single-chain polypeptide with a molecular weight of 110,000. It is present in plasma in an average concentration of 25 µg/ml. The activated form of C2 combines with the activated C4 to form a complex with C3-conv |
| Formel                 | Delipidated, heat inactivated, lyophilized, stable IgG fraction. No preservative added. The IgG concentration is 4 mg/ml in PBS-EDTA. No foreign proteins added.   |
| Anwendungsbeschreibung | Precipitation assays. In immunoelectrophoresis in agarose-plates use 2 µl human plasma or equivalent against 120 µl antiserum. In double radial immunodiffusion use a rosette arrangement with 10 µl antiserum in 3 mm diameter center well and 2 µl plasma sa |