

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Rabbit anti Human lysozyme, conjugated with Horseradish peroxidase, Clone: [Polyclonal], HRP, Monoclonal NMB-RAHU/LYS/PO

| | |
|--------------------------|---|
| Artikelname | Rabbit anti Human lysozyme, conjugated with Horseradish peroxidase, Clone: [Polyclonal], HRP, Monoclonal |
| Artikelnummer | NMB-RAHU/LYS/PO |
| Hersteller Artikelnummer | RAHu/Lys/PO |
| Alternativnummer | NMB-RAHU/LYS/PO |
| Hersteller | NordicMubio |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Spezies Reaktivität | Human |
| Immunogen | Highly purified recombinant human lysozyme was used as immunogen. |
| Konjugation | HRP |
| Produktbeschreibung | The reactivity of the antiserum is restricted to lysozyme. In immunoelectrophoresis and radial immunodiffusion (Ouchterlony), using various antiserum concentrations against normal human milk and tears a single precipitin line is obtained which shows ... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [Polyclonal] |
| UniProt | P61626 |

| | |
|------------------------|--|
| Puffer | Horseradish peroxidase-coupled purified hyperimmune IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2) No preservative added, as it may interfere with the antibody activity. |
| Quelle | Lysozyme, a bacteriolytic enzyme, has a molecular weight of about 15,000. Its concentration in plasma is low, in milk the concentration is reported to be between 5 and 220 mg/ml. Also tears contain a considerable amount of lysozyme. Highly purified recom |
| Formel | Horseradish peroxidase-coupled purified hyperimmune IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2) No preservative added, as it may interfere with the antibody activity. |
| Anwendungsbeschreibung | ELISA, Immunocytochemistry, Immunohistochemistry (paraffin), Dot blot, Immunoblotting. The lyophilized conjugate is shipped at ambient temperature and may be stored at +4C, prolonged storage at or below -20C. It is reconstituted by adding 1 ml sterile di s |