

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Rabbit anti Human platelet factor 4, Clone: [Polyclonal], Monoclonal NMB-RAHU/PF4

|                          |  |
|--------------------------|--|
| Artikelname              | Rabbit anti Human platelet factor 4, Clone: [Polyclonal], Monoclonal   |
| Artikelnummer            | NMB-RAHU/PF4   |
| Hersteller Artikelnummer | RAHu/PF4   |
| Alternativnummer         | NMB-RAHU/PF4   |
| Hersteller               | NordicMubio  |
| Wirt                     | Rabbit   |
| Kategorie                | Antikörper   |
| Produktbeschreibung      | The reactivity of the antiserum is restricted to platelet factor 4 as tested in immunoelectrophoresis and radial immunodiffusion. A characteristic single precipitin line is obtained with normal serum which shows a reaction of identity with the precip...  |
| Klonalität               | Monoclonal   |
| Klon-Bezeichnung         | [Polyclonal]   |
| UniProt                  | <a href="#">P02776</a>   |
| Puffer                   | Delipidated, heat inactivated, lyophilized, stable whole antiserum Sodium azide. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added. Reconstitute the lyophilized antiseru |
| Quelle                   | Platelet factor 4 is a low molecular weight (MW 7,800) heat stable protein, secreted by alpha granules of the platelets. Purified platelet factor 4 isolated from pooled human serum. Freund's complete adjuvant is used in the first step of the immunization |

|                        |  |
|------------------------|--|
| Formel                 | Delipidated, heat inactivated, lyophilized, stable whole antiserum Sodium azide. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added.                                       |
| Anwendungsbeschreibung | Precipitation assays. In immunoelectrophoresis use 2 µl serum or equivalent against 120 µl antiserum. In double radial immunodiffusion use a rosette arrangement with 10 µl antiserum in 3 mm diameter centre well and 2 µl serum samples (neat and serially d |