

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Rabbit anti Mouse alpha-2 macroglobulin, Clone: [Polyclonal], Monoclonal NMB-RAM/A2M

|                          |   |
|--------------------------|---|
| Artikelname              | Rabbit anti Mouse alpha-2 macroglobulin, Clone: [Polyclonal], Monoclonal  |
| Artikelnummer            | NMB-RAM/A2M   |
| Hersteller Artikelnummer | RAM/A2M   |
| Alternativnummer         | NMB-RAM/A2M   |
| Hersteller               | NordicMubio   |
| Wirt                     | Rabbit  |
| Kategorie                | Antikörper  |
| Produktbeschreibung      | The reactivity of the antiserum is restricted to alpha-2 macroglobulin as tested in immunoelectrophoresis and radial immunodiffusion. A characteristic single precipitin line is obtained with normal serum which shows a reaction of identity with the pr... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [Polyclonal]  |
| Puffer                   | Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added. Reconstitute the lyophilized |
| Quelle                   | Alpha-2 macroglobulin is the major protein in the alpha-2 region of mouse serum after separation by electrophoresis. It is a large polymer and a strong antigen. It forms about one-third in the total of the alpha-2 globulins. The protein is isolated by a |

|                        |  |
|------------------------|--|
| Formel                 | Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added.                               |
| Anwendungsbeschreibung | Precipitation assays. In immunoelectrophoresis use 2 µl serum or equivalent against 120 µl antiserum. In double radial immunodiffusion use a rosette arrangement with 10 µl antiserum in 3 mm diameter centre well and 2 µl serum samples (neat and serially d |