

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Rabbit anti Mouse IgE (Fc specific), Clone: [Polyclonal], Monoclonal NMB-RAM/IGE(FC)

| | |
|--------------------------|--|
| Artikelname | Rabbit anti Mouse IgE (Fc specific), Clone: [Polyclonal], Monoclonal |
| Artikelnummer | NMB-RAM/IGE(FC) |
| Hersteller Artikelnummer | RAM/IgE(Fc) |
| Alternativnummer | NMB-RAM/IGE(FC) |
| Hersteller | NordicMubio |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Produktbeschreibung | The reactivity of the antiserum is directed to the Fc subunits of IgE as tested against mouse sera and a panel of purified homogenous immunoglobulins. It does not react with IgG, IgG/Fab fragments, IgM and IgA or any non-Ig protein in mouse serum, as... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [Polyclonal] |
| Puffer | Delipidated, heat inactivated, lyophilized stable whole serum No preservative added. Total protein and IgG concentration in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added. Reconstitute the lyophilized antis |
| Quelle | Pools of purified homogenous IgE isolated from pooled mouse serum. Freund's complete adjuvant is used in the first step of the immunization procedure. |

| | |
|------------------------|---|
| Formel | Delipidated, heat inactivated, lyophilized stable whole serum No preservative added. Total protein and IgG concentration in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added. |
| Anwendungsbeschreibung | Precipitation assays. In immunoelectrophoresis use 2 μ l or equivalent against 120 μ l antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 μ l antiserum in a 3 mm diameter centre well and 2 μ l serum samples (neat and |