

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Rabbit anti Swine secretory IgA, Clone: [Polyclonal], Monoclonal NMB-RASW/SIGA

| | |
|--------------------------|--|
| Artikelname | Rabbit anti Swine secretory IgA, Clone: [Polyclonal], Monoclonal |
| Artikelnummer | NMB-RASW/SIGA |
| Hersteller Artikelnummer | RASw/sIgA |
| Alternativnummer | NMB-RASW/SIGA |
| Hersteller | NordicMubio |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Produktbeschreibung | The reactivity of the antiserum is restricted to the Fc part of the IgA molecule and the surface and hidden determinants of the secretory component. In immunoelectrophoresis and radial immunodiffusion, using various antiserum concentrations against n... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [Polyclonal] |
| Puffer | Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added. Reconstitute the lyophilize |
| Quelle | Highly purified secretory IgA isolated from pooled swine milk. Freund's complete adjuvant is used in the first step of the immunization procedure. |

| | |
|------------------------|---|
| Formel | Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added. |
| Anwendungsbeschreibung | Precipitation assays. In immunoelectrophoresis use 2 µl serum or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in 3 mm diameter center well and 2 µl serum samples (neat |