

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

ABO Antibody (Blood Group Antigen A), Clone: [HE-14], Mouse, Monoclonal NSJ-V2549SAF-100UG

Artikelname	ABO Antibody (Blood Group Antigen A), Clone: [HE-14], Mouse, Monoclonal
Artikelnummer	NSJ-V2549SAF-100UG
Hersteller Artikelnummer	V2549SAF-100UG
Alternativnummer	NSJ-V2549SAF-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IF, IH
Spezies Reaktivität	Human
Immunogen	A mixture of erythrocytes of blood group A1 and glycoprotein fraction isolated from the saliva of secretors with blood group A was used as the immunogen for the ABO antibody.
Produktbeschreibung	This antibody recognizes human blood group A (monofucosyl and difucosyl A antigens with chain types 1, 2, 3, 4, 5, 6) and Forssmann antigen. It is also reactive with the immuno-dominant A trisaccharide. Blood group antigen expression in human colon c...
Klonalität	Monoclonal
Konzentration	1 mg/ml in 1X PBS, BSA free, sodium azide free
Klon-Bezeichnung	[HE-14]

Isotyp	Mouse IgM, kappa
UniProt	P16442
Puffer	1 mg/ml in 1X PBS, BSA free, sodium azide free
Reinheit	PEG precipitation
Formulierung	1 mg/ml in 1X PBS, BSA free, sodium azide free
Target-Kategorie	ABO
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistology (formalin-fixed): 1-2ug/ml, Immunofluorescence: 2-4ug/ml
Anwendungsbeschreibung	Optimal dilution of the ABO antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes