

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD13 Antibody / ANPEP, Clone: [APN/514], Mouse, Monoclonal NSJ-V2557-20UG

| | |
|--------------------------|---|
| Artikelname | CD13 Antibody / ANPEP, Clone: [APN/514], Mouse, Monoclonal |
| Artikelnummer | NSJ-V2557-20UG |
| Hersteller Artikelnummer | V2557-20UG |
| Alternativnummer | NSJ-V2557-20UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | ELISA |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human CD13 was used as the immunogen for the CD13 antibody. |
| Produktbeschreibung | This mAb recognizes an extracellular epitope of an integral membrane glycoprotein of 150kDa, identified as CD13. This antigen is present on most cells of myeloid origin including granulocytes, monocytes, mast cells, and GM-progenitor cells. It is als... |
| Klonalität | Monoclonal |
| Konzentration | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide |
| Klon-Bezeichnung | [APN/514] |
| Isotyp | Mouse IgG1, kappa |

| | |
|------------------------|--|
| UniProt | P15144 |
| Puffer | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide |
| Reinheit | Protein G affinity chromatography |
| Formulierung | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide |
| Target-Kategorie | CD13 |
| Antibody Type | Primary Antibody |
| Application Verdünnung | ELISA (order BSA/sodium azide-free format for coating) |
| Anwendungsbeschreibung | Optimal dilution of the CD13 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min |