

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### HLA-DPB1 Antibody (MHC II), Clone: [BraFB6], Mouse, Monoclonal NSJ-V2578-100UG

Artikelname	HLA-DPB1 Antibody (MHC II), Clone: [BraFB6], Mouse, Monoclonal
Artikelnummer	NSJ-V2578-100UG
Hersteller Artikelnummer	V2578-100UG
Alternativnummer	NSJ-V2578-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	ELISA, FACS, IF
Spezies Reaktivität	Human
Immunogen	Non-T, non-B human acute lymphoblastic leukemia REH6 cell line was used as the immunogen for the HLA-DPB1 antibody.
Produktbeschreibung	Recognizes a non-polymorphic determinant of DP-MHC class II. MHC class II antigens are transmembrane glycoproteins of non-covalently linked alpha (33-35kDa) and beta (27-30kDa) chains. It reportedly reacts with B- & non-T, non-B cell lines but not wi...
Klonalität	Monoclonal
Konzentration	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Klon-Bezeichnung	[BraFB6]
Isotyp	Mouse IgG2b, kappa

UniProt	<a href="#">P04440</a>
Puffer	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Reinheit	Protein G affinity chromatography
Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Target-Kategorie	HLA-DPB1
Antibody Type	Primary Antibody
Application Verdünnung	ELISA: 2-4ug/ml for coating (order BSA/sodium azide-free format),Flow cytometry: 1-2ug/million cells,Immunofluorescence: 1-2ug/ml
Anwendungsbeschreibung	Optimal dilution of the HLA-DPB1 antibody should be determined by the researcher.1. Staining of formalin/paraffin tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.