

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

TLR4 Antibody, Clone: [TLR4/230], Mouse, Monoclonal NSJ-V2897-100UG

| | |
|--------------------------|---|
| Artikelname | TLR4 Antibody, Clone: [TLR4/230], Mouse, Monoclonal |
| Artikelnummer | NSJ-V2897-100UG |
| Hersteller Artikelnummer | V2897-100UG |
| Alternativnummer | NSJ-V2897-100UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | Functional Studies |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human protein was used as the immunogen for the TLR4 antibody. |
| Produktbeschreibung | This mAb reacts with human Toll-like receptor 2 (TLR4). It is a member of the Toll-like receptor (TLR) family, which plays a fundamental role in pathogen recognition and activation of innate immunity. TLRs are highly conserved from Drosophila to huma... |
| Klonalität | Monoclonal |
| Konzentration | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide |
| Klon-Bezeichnung | [TLR4/230] |
| Isotyp | Mouse IgG2a, kappa |

| | |
|------------------------|---|
| UniProt | O00206 |
| Puffer | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide |
| Reinheit | Protein G affinity chromatography |
| Formulierung | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide |
| Target-Kategorie | TLR4 |
| Antibody Type | Primary Antibody |
| Application Verdünnung | Functional studies (order BSA/sodium azide-free format) |
| Anwendungsbeschreibung | Optimal dilution of the TLR2 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min. |