

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### HDAC1 Monoclonal Antibody [10E2] - Unconjugated / 0.1ml, IgG1, Mouse IMQ-IQ555

|                          |  |
|--------------------------|--|
| Artikelname              | HDAC1 Monoclonal Antibody [10E2] - Unconjugated / 0.1ml, IgG1, Mouse   |
| Artikelnummer            | IMQ-IQ555  |
| Hersteller Artikelnummer | IQ555  |
| Alternativnummer         | IMQ-IQ555-0.1  |
| Hersteller               | ImmuQuest  |
| Wirt                     | Mouse  |
| Kategorie                | Antikörper   |
| Applikation              | ChIP, FC, ICC, IHC-P, IP, WB   |
| Spezies Reaktivität      | Human, Mouse, Rat  |
| Immunogen                | Synthetic peptide: EEKPEAKGVKKEEVKLA, corresponding to C terminal amino acids 467-482 of Human and mouse HDAC1 |
| Produktbeschreibung      | Mouse Anti-HDAC1 [10E2]...   |
| Klonalität               | Monoclonal   |
| Konzentration            | 1mg/ml   |
| Klon-Bezeichnung         | [10E2]   |
| Isotyp                   | IgG1   |
| NCBI                     | <a href="#">3065</a>   |

|                        |   |
|------------------------|---|
| Pubmed                 | <a href=https://pubmed.ncbi.nlm.nih.gov/12032080 target=_blank rel=nofollow>12032080  |
| UniProt                | <a href="#">Q13547</a>  |
| Puffer                 | Purified antibody (from supernatant) containing PBS + 0.1% sodium azide   |
| Reinheit               | Protein G   |
| Formulierung           | Purified antibody (from supernatant) containing PBS + 0.1% sodium azide   |
| Target-Kategorie       | HDAC1   |
| Antibody Type          | Monoclonal Antibody   |
| Application Verdünnung | Optimal antibody dilution should be determined by titration, however as a guideline try, A WB: 1/500 - 1/4000. Detects a band of approximately 57 kDa (predicted molecular weight: 55 kDa). |