

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

p27Kip1 (Mitotic Inhibitor/Suppressor Protein) Antibody, IgG1, Clone: [SX53G8], Mouse, Monoclonal NBT-1027-MSM1-P0

| | |
|--------------------------|---|
| Artikelname | p27Kip1 (Mitotic Inhibitor/Suppressor Protein) Antibody, IgG1, Clone: [SX53G8], Mouse, Monoclonal |
| Artikelnummer | NBT-1027-MSM1-P0 |
| Hersteller Artikelnummer | 1027-MSM1-P0 |
| Alternativnummer | NBT-1027-MSM1-P0-20,NBT-1027-MSM1-P0-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC |
| Spezies Reaktivität | Human, Monkey, Mouse, Rat |
| Immunogen | Purified GST-p27 fusion protein of human origin |
| Produktbeschreibung | This MAb recognizes a 27kDa protein, identified as the p27Kip1, a cell cycle regulatory mitotic inhibitor. It is highly specific and shows no cross-reaction with other related mitotic inhibitors. In Western blotting of cell lysates from 7 human breas... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [SX53G8] |
| Molekulargewicht | 25-26kDa |
| Isotyp | IgG1 |

| | |
|------------------------|--|
| NCBI | 1027 |
| UniProt | P46527 |
| Formulierung | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (0.25-0.5ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 |