

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cystatin A Antibody, IgG2c, Clone: [CPTC-CSTA-1], Mouse, Monoclonal NBT-1475-MSM1-P1ABX

| | |
|--------------------------|---|
| Artikelname | Cystatin A Antibody, IgG2c, Clone: [CPTC-CSTA-1], Mouse, Monoclonal |
| Artikelnummer | NBT-1475-MSM1-P1ABX |
| Hersteller Artikelnummer | 1475-MSM1-P1ABX |
| Alternativnummer | NBT-1475-MSM1-P1ABX-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human CSTA protein |
| Produktbeschreibung | The cystatin superfamily encompasses proteins that contain multiple cystatin-like sequences. Some of the members are active cysteine protease inhibitors, while others have lost or perhaps never acquired this inhibitory activity. There are three inhib... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [CPTC-CSTA-1] |
| Molekulargewicht | 11kDa |
| Isotyp | IgG2c |

| | |
|------------------------|---|
| NCBI | 1475 |
| UniProt | P01040 |
| Formulierung | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Western Blot (2-4ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes |