

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Erythropoietin (EPO) (Marker of Placentation Disorders) Antibody, IgG1, Clone: [EPO/1368], Mouse, Monoclonal NBT-2056-MSM2-P1ABX

| | |
|--------------------------|---|
| Artikelname | Erythropoietin (EPO) (Marker of Placentation Disorders) Antibody, IgG1, Clone: [EPO/1368], Mouse, Monoclonal |
| Artikelnummer | NBT-2056-MSM2-P1ABX |
| Hersteller Artikelnummer | 2056-MSM2-P1ABX |
| Alternativnummer | NBT-2056-MSM2-P1ABX-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | ELISA, IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant fragment (around aa 28-162) of human EPO protein (exact sequence is proprietary) |
| Produktbeschreibung | Recognizes a protein of about 37kDa, which is identified as Erythropoietin (EPO). Erythropoietin is a secreted, glycosylated cytokine hormone composed of four alpha helical bundles. It is the primary factor responsible for regulating erythropoiesis d... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [EPO/1368] |
| Molekulargewicht | 37kDa |

| | |
|------------------------|--|
| Isotyp | IgG1 |
| NCBI | 2056 |
| UniProt | P01588 |
| Formulierung | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | ELISA (Use Ab at 2-4ug/ml for coating) (Order Ab without BSA), Western Blot (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (2-4ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9 |