

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

AKT1 (Prognostic Marker for Neuroendocrine Tumors), IgG1, Clone: [rAKT1/2491], Mouse, Monoclonal NBT-207-MSM5-P0

| | |
|--------------------------|---|
| Artikelname | AKT1 (Prognostic Marker for Neuroendocrine Tumors), IgG1, Clone: [rAKT1/2491], Mouse, Monoclonal |
| Artikelnummer | NBT-207-MSM5-P0 |
| Hersteller Artikelnummer | 207-MSM5-P0 |
| Alternativnummer | NBT-207-MSM5-P0-20,NBT-207-MSM5-P0-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant fragment of human AKT1 protein (around aa 85-189) (exact sequence is proprietary) |
| Produktbeschreibung | Recognizes a protein of 62kDa, which is identified as AKT1The serine/threonine kinase Akt family contains several members, including Akt1 (also designated PKB or RacPK), Akt2 (also designated PKB or RacPK-) and Akt 3 (also designated PKB or thymoma v... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [rAKT1/2491] |
| Molekulargewicht | 62kDa |

| | |
|------------------------|--|
| Isotyp | IgG1 |
| NCBI | 207 |
| UniProt | P31749 |
| Formulierung | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 1mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Anwendungsbeschreibung | Western Blot (1-2ug/ml), ,Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 min at RT),(Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate Buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes),Optim |