

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Fibroblast Activation Protein Alpha / FAP-1 Antibody, IgG2c, Clone: [FAP/4851], Mouse, Monoclonal NBT-2191-MSM1-P0**

|                          |   |
|--------------------------|---|
| Artikelname              | Fibroblast Activation Protein Alpha / FAP-1 Antibody, IgG2c, Clone: [FAP/4851], Mouse, Monoclonal   |
| Artikelnummer            | NBT-2191-MSM1-P0  |
| Hersteller Artikelnummer | 2191-MSM1-P0  |
| Alternativnummer         | NBT-2191-MSM1-P0-20,NBT-2191-MSM1-P0-100  |
| Hersteller               | NeoBiotechnologies  |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant fragment (around aa 1-200) of human FAP protein (exact sequence is proprietary)   |
| Produktbeschreibung      | FAP (fibroblast activation protein) is a cell surface glycoprotein and serine protease that is expressed primarily in fetal mesenchymal tissues and epithelial cancer fibroblasts. In cancer, FAP functions to promote cellular proliferation. In embryoni... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [FAP/4851]  |
| Molekulargewicht         | 88kDa   |

|                        |   |
|------------------------|---|
| Isotyp                 | IgG2c   |
| NCBI                   | <a href="#">2191</a>  |
| UniProt                | <a href="#">Q12884</a>  |
| Formulierung           | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.  |
| Antibody Type          | Monoclonal Antibody   |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal diluti |