

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Fibrillin-1 (FBN1) (Marfan Syndrome Marker) Antibody, IgG1, Clone: [FBN1/2191], Mouse, Monoclonal NBT-2200-MSM1-P0**

|                          |   |
|--------------------------|---|
| Artikelname              | Fibrillin-1 (FBN1) (Marfan Syndrome Marker) Antibody, IgG1, Clone: [FBN1/2191], Mouse, Monoclonal   |
| Artikelnummer            | NBT-2200-MSM1-P0  |
| Hersteller Artikelnummer | 2200-MSM1-P0  |
| Alternativnummer         | NBT-2200-MSM1-P0-20,NBT-2200-MSM1-P0-100  |
| Hersteller               | NeoBiotechnologies  |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant human Fibrillin 1 fragment around aa 330-476 (Exact sequence is proprietary)  |
| Produktbeschreibung      | Fibrillin-1 is an extracellular matrix glycoprotein that serves as a structural component of calcium-binding microfibrils. These microfibrils provide force-bearing structural support in elastic and non-elastic connective tissue throughout the body.Mu... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [FBN1/2191]   |
| Molekulargewicht         | 350kDa  |

|                        |   |
|------------------------|---|
| Isotyp                 | IgG1  |
| NCBI                   | <a href="#">2200</a>  |
| UniProt                | <a href="#">P35555</a>  |
| Formulierung           | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.  |
| Antibody Type          | Monoclonal Antibody   |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal diluti |