

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **GFAP (Astrocyte & Neural Stem Cell Marker) Antibody, IgG1, Clone: [SPM507], Mouse, Monoclonal NBT-2670-MSM1XX-P1ABX**

|                          |   |
|--------------------------|---|
| Artikelname              | GFAP (Astrocyte & Neural Stem Cell Marker) Antibody, IgG1, Clone: [SPM507], Mouse, Monoclonal   |
| Artikelnummer            | NBT-2670-MSM1XX-P1ABX   |
| Hersteller Artikelnummer | 2670-MSM1XX-P1ABX   |
| Alternativnummer         | NBT-2670-MSM1XX-P1ABX-100   |
| Hersteller               | NeoBiotechnologies  |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | FC, IF, IHC, WB   |
| Spezies Reaktivität      | Bovine, Gallus, Human, Mouse, Porcine, Rabbit, Rat  |
| Immunogen                | GFAP isolated from pig spinal cord  |
| Produktbeschreibung      | This MAb recognizes a protein of ~50kDa which is identified as Glial Fibrillary Acidic Protein (GFAP). It shows no cross-reaction with other intermediate filament proteins. GFAP is specifically found in astroglia. GFAP is a very popular marker for lo... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [SPM507]  |
| Molekulargewicht         | ~50kDa  |
| Isotyp                   | IgG1  |

|                        |   |
|------------------------|---|
| NCBI                   | <a href="#">2670</a>  |
| UniProt                | <a href="#">P14136</a>  |
| Formulierung           | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.  |
| Antibody Type          | Monoclonal Antibody   |
| Anwendungsbeschreibung | Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Western Blot (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM |