

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Apolipoprotein E / APOE Antibody, IgG1, Clone: [APOE/3671], Mouse, Monoclonal NBT-348-MSM1-P0

| | |
|--------------------------|---|
| Artikelname | Apolipoprotein E / APOE Antibody, IgG1, Clone: [APOE/3671], Mouse, Monoclonal |
| Artikelnummer | NBT-348-MSM1-P0 |
| Hersteller Artikelnummer | 348-MSM1-P0 |
| Alternativnummer | NBT-348-MSM1-P0-20,NBT-348-MSM1-P0-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Human recombinant APOE protein fragment (around aa19-119) (exact sequence is proprietary) |
| Produktbeschreibung | In skeletal muscle, AQP4 (aquaporin 4 also known as mercurial insensitive water channel), localizes to the sarcolemma of fast-twitch muscle fibers. Aquaporins (AQPs) are a large family of integral membrane water transport channel proteins that facili... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [APOE/3671] |
| Molekulargewicht | 36kDa |

| | |
|------------------------|---|
| Isotyp | IgG1 |
| NCBI | 348 |
| UniProt | P02649 |
| Formulierung | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal diluti |