

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

ARF1 (Golgi Apparatus Marker) Antibody, IgG2a, Clone: [1A9/5], Mouse, Monoclonal NBT-375-MSM1-P0

| | |
|--------------------------|---|
| Artikelname | ARF1 (Golgi Apparatus Marker) Antibody, IgG2a, Clone: [1A9/5], Mouse, Monoclonal |
| Artikelnummer | NBT-375-MSM1-P0 |
| Hersteller Artikelnummer | 375-MSM1-P0 |
| Alternativnummer | NBT-375-MSM1-P0-20,NBT-375-MSM1-P0-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Human, Mouse, Rat |
| Immunogen | C-terminal human ARF1 peptide SNQLRNQ |
| Produktbeschreibung | The ADP-ribosylation factor (ARF) family comprises a group of structurally and functionally conserved proteins, which are members of the Ras superfamily of regulatory GTP-binding proteins. The ARF family is divided functionally into the ARF and the A... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [1A9/5] |
| Molekulargewicht | 20kDa |
| Isotyp | IgG2a |

| | |
|------------------------|---|
| NCBI | 375 |
| UniProt | P84077 |
| Formulierung | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Western Blot (1-2ug/ml),Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes) |