

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin 5 (KRT5) (Basal, Myoepithelial & Mesothelial Cell Marker) Antibody, IgG1, Clone: [KRT5/2080], Mouse, Monoclonal NBT-3852-MSM1-CS0

| | |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname | Cytokeratin 5 (KRT5) (Basal, Myoepithelial & Mesothelial Cell Marker) Antibody, IgG1, Clone: [KRT5/2080], Mouse, Monoclonal |
| Artikelnummer | NBT-3852-MSM1-CS0 |
| Hersteller Artikelnummer | 3852-MSM1-CS0 |
| Alternativnummer | NBT-3852-MSM1-CS0-0.1,NBT-3852-MSM1-CS0-0.5,NBT-3852-MSM1-CS0-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human Cytokeratin 5 (KRT5) protein fragment (around aa 318-491) (exact sequence is proprietary) |
| Produktbeschreibung | This MAb recognizes a protein of 58kDa, which is identified as Cytokeratin 5 (KRT5). This type II cytokeratin is specifically expressed in the basal layer of the epidermis with family member KRT14. Antibodies to KRT5 identify basal cells of squamous ... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [KRT5/2080] |
| Molekulargewicht | 58kDa |

| | |
|------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Isotyp | IgG1 |
| NCBI | 3852 |
| UniProt | P13647 |
| Formulierung | Tissue culture supernatant with 0.05% Azide. Contact us if you require it in a different format. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1:50-1:100 for 30 minutes at RT) (Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilutio |