

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin 15 (Esophageal Squamous Cell Carcinoma Marker) Antibody, IgG2a, Clone: [SPM190], Mouse, Monoclonal NBT-3866-MSM1X-P1ABX

| | |
|--------------------------|---|
| Artikelname | Cytokeratin 15 (Esophageal Squamous Cell Carcinoma Marker) Antibody, IgG2a, Clone: [SPM190], Mouse, Monoclonal |
| Artikelnummer | NBT-3866-MSM1X-P1ABX |
| Hersteller Artikelnummer | 3866-MSM1X-P1ABX |
| Alternativnummer | NBT-3866-MSM1X-P1ABX-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Bovine, Human, Rat |
| Immunogen | A 17 amino acids synthetic peptide from the C- terminal of Human Cytokeratin 15 |
| Produktbeschreibung | Keratin 15 is a type I keratin which is expressed only in basal keratinocytes in stratified epithelia and does not appear to have a natural type II expression partner. Keratin 15 is down regulated in activated keratinocytes. Cytokeratin 15 is a speci... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [SPM190] |
| Molekulargewicht | 52kDa |

| | |
|------------------------|---|
| Isotyp | IgG2a |
| NCBI | 3866 |
| UniProt | P19012 |
| Formulierung | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Immunohistochemistry (Frozen & formalin-fixed) (1-2ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilut |