

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Recombinant Cytokeratin 16 (KRT16) (Suprabasal Keratinocyte Marker) Antibody, IgG1, Clone: [rKRT16/1714], Mouse, Monoclonal NBT-3868-MSM4-P1ABX**

|                          |   |
|--------------------------|---|
| Artikelname              | Recombinant Cytokeratin 16 (KRT16) (Suprabasal Keratinocyte Marker) Antibody, IgG1, Clone: [rKRT16/1714], Mouse, Monoclonal   |
| Artikelnummer            | NBT-3868-MSM4-P1ABX   |
| Hersteller Artikelnummer | 3868-MSM4-P1ABX   |
| Alternativnummer         | NBT-3868-MSM4-P1ABX-100   |
| Hersteller               | NeoBiotechnologies  |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant fragment from the C-terminal of human Cytokeratin 16  |
| Produktbeschreibung      | Cytokeratins are a family of intermediate filament proteins that assemble into filaments through forming heterodimers of one type I Cytokeratins (Cytokeratins 9 to 23) and one type II Cytokeratins (keratins 1 to 8). The cytokeratin proteins play a cr... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [rKRT16/1714]   |
| Molekulargewicht         | 48kDa   |
| Isotyp                   | IgG1  |

|                        |  |
|------------------------|--|
| NCBI                   | <a href="#">3868</a>   |
| UniProt                | <a href="#">P08779</a>   |
| Formulierung           | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.   |
| Antibody Type          | Recombinant Monoclonal Antibody  |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilution |