

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

MDM2 Antibody, IgG1, Clone: [MDM2/3277], Mouse, Monoclonal NBT-4193-MSM6-P1ABX

| | |
|--------------------------|---|
| Artikelname | MDM2 Antibody, IgG1, Clone: [MDM2/3277], Mouse, Monoclonal |
| Artikelnummer | NBT-4193-MSM6-P1ABX |
| Hersteller Artikelnummer | 4193-MSM6-P1ABX |
| Alternativnummer | NBT-4193-MSM6-P1ABX-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human MDM2 protein fragment (around aa 126-254) (exact sequence is proprietary) |
| Produktbeschreibung | MDM2 is a nuclear phosphoprotein that binds and inhibits transactivation by tumor protein p53. It can promote tumor formation by targeting tumor suppressor proteins, such as p53, for proteasomal degradation. Overexpression of MDM2 can result in exces... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [MDM2/3277] |
| Molekulargewicht | ~90kDa |
| Isotyp | IgG1 |

| | |
|------------------------|---|
| NCBI | 4193 |
| UniProt | Q00987 |
| Formulierung | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal diluti |