

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

MUC1 / CA15-3 / EMA / CD227 (Epithelial Marker) Antibody, IgG1, Clone: [MUC1/967], Mouse, Monoclonal NBT-4582-MSM17-P1ABX

| | |
|--------------------------|--|
| Artikelname | MUC1 / CA15-3 / EMA / CD227 (Epithelial Marker) Antibody, IgG1, Clone: [MUC1/967], Mouse, Monoclonal |
| Artikelnummer | NBT-4582-MSM17-P1ABX |
| Hersteller Artikelnummer | 4582-MSM17-P1ABX |
| Alternativnummer | NBT-4582-MSM17-P1ABX-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF, WB |
| Spezies Reaktivität | Human |
| Immunogen | Human milk-fat globule membranes (HMFGM) |
| Produktbeschreibung | This MAb recognizes full-length MUC1 in a glycosylation-independent manner and can bind to the fully glycosylated protein. The dominant epitope of this MAb is APDTR in the VNTR region. It reacts with the core peptide of the MUC1 protein, which is a me... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [MUC1/967] |
| Molekulargewicht | 265-400kDa |
| Isotyp | IgG1 |

| | |
|------------------------|---|
| NCBI | 4582 |
| UniProt | P15941 |
| Formulierung | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Western Blot (1-2ug/ml), Flow Cytometry (0.5-1ug/million cells), Immunofluorescence (1-2ug/ml), Immunohistology (Formalin-paraffin) (0.25-0.5ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate |