

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

MUC1 / CA15-3 / EMA / CD227 (Epithelial Marker) Antibody, Clone: [GP1.4 + E29], Mouse, Monoclonal NBT-4582-MSM3-P1ABX

| | |
|--------------------------|---|
| Artikelname | MUC1 / CA15-3 / EMA / CD227 (Epithelial Marker) Antibody, Clone: [GP1.4 + E29], Mouse, Monoclonal |
| Artikelnummer | NBT-4582-MSM3-P1ABX |
| Hersteller Artikelnummer | 4582-MSM3-P1ABX |
| Alternativnummer | NBT-4582-MSM3-P1ABX-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Human milk fat globule membranes (GP1.4), Delipidated extract of human milk fat globule membranes (E29) |
| Produktbeschreibung | In Western blotting, it recognizes proteins in MW range of 265-400kDa, identified as different glycoforms of EMA. The alpha subunit has cell adhesive properties. It can act both as an adhesion and an anti-adhesion protein. EMA may provide a protectiv... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [GP1.4 + E29] |
| Molekulargewicht | 265-400kDa |

| | |
|------------------------|---|
| NCBI | 4582 |
| UniProt | P15941 |
| Formulierung | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilution for a |