

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Recombinant CD31 / PECAM-1 (Endothelial Cell Marker) Antibody, IgG2a, Clone: [r6439], Mouse, Monoclonal NBT-5175-MSM36-P0

| | |
|--------------------------|---|
| Artikelname | Recombinant CD31 / PECAM-1 (Endothelial Cell Marker) Antibody, IgG2a, Clone: [r6439], Mouse, Monoclonal |
| Artikelnummer | NBT-5175-MSM36-P0 |
| Hersteller Artikelnummer | 5175-MSM36-P0 |
| Alternativnummer | NBT-5175-MSM36-P0-20,NBT-5175-MSM36-P0-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human CD31 protein |
| Produktbeschreibung | CD31 (PECAM-1) is a transmembrane glycoprotein member of the immunoglobulin supergene family of adhesion molecules. CD31 is expressed by stem cells of the hematopoietic system and is primarily used to identify and concentrate these cells for experime... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [r6439] |
| Molekulargewicht | ~100kDa (endothelium) and ~130kDa (platelets) |
| Isotyp | IgG2a |

| | |
|------------------------|--|
| NCBI | 5175 |
| UniProt | P16284 |
| Formulierung | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Anwendungsbeschreibung | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes),Optimal dilutio |