

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Aciculin / Phosphoglucomutase 5 (PGM5) Antibody, IgG2a, Clone: [14F8/F8], Mouse, Monoclonal NBT-5239-MSM1-P1ABX**

|                          |   |
|--------------------------|---|
| Artikelname              | Aciculin / Phosphoglucomutase 5 (PGM5) Antibody, IgG2a, Clone: [14F8/F8], Mouse, Monoclonal   |
| Artikelnummer            | NBT-5239-MSM1-P1ABX   |
| Hersteller Artikelnummer | 5239-MSM1-P1ABX   |
| Alternativnummer         | NBT-5239-MSM1-P1ABX-100   |
| Hersteller               | NeoBiotechnologies  |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | ELISA, IF, IHC, WB  |
| Spezies Reaktivität      | Bovine, Human, Mouse, Rabbit, Rat   |
| Immunogen                | Cytoskeletal fraction of the human myometrium tissue extract  |
| Produktbeschreibung      | Aciculin, also known as PGM5 (phosphoglucomutase 5) or PGMRP, is a 567 amino acid protein that belongs to the phosphoglucomutase family of phosphotransferases, which play an important role in the interconversion of glucose-1-phosphate and glucose-6-p... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [14F8/F8]   |
| Molekulargewicht         | 56kDa   |
| Isotyp                   | IgG2a   |

|                        |  |
|------------------------|--|
| NCBI                   | <a href="#">5239</a>   |
| UniProt                | <a href="#">Q15124</a>   |
| Formulierung           | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.   |
| Antibody Type          | Monoclonal Antibody  |
| Anwendungsbeschreibung | ELISA (For coating, order antibody without BSA), Immunofluorescence (1-2ug/ml), Western Blot (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT), (Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tri |