

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

SERPINB5 / Maspin Antibody, IgG1, Clone: [SERPINB5/4972], Mouse, Monoclonal NBT-5268-MSM2-P1ABX

| | |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname | SERPINB5 / Maspin Antibody, IgG1, Clone: [SERPINB5/4972], Mouse, Monoclonal |
| Artikelnummer | NBT-5268-MSM2-P1ABX |
| Hersteller Artikelnummer | 5268-MSM2-P1ABX |
| Alternativnummer | NBT-5268-MSM2-P1ABX-100 |
| Hersteller | NeoBiotechnologies |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant fragment of human SERPINB5 protein (around aa1-200) (exact sequence is proprietary) |
| Produktbeschreibung | Maspin is structurally a serine protease inhibitor (serpin) that was initially isolated from normal human mammary epithelial cells. Serpins are a family of proteins that inhibit Chymotrypsin-like serine proteinases. Serpins control activated proteina... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [SERPINB5/4972] |
| Molekulargewicht | 42kDa |

| | |
|------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Isotyp | IgG1 |
| NCBI | 5268 |
| UniProt | P36952 |
| Formulierung | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Anwendungsbeschreibung | Western Blot (1-2ug/ml), ,Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for |